

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

1980

Symbiotic Nitrogen Fixation by Legumes Requires Extra Phosphorus

Kenneth G. Cassman

University of Nebraska-Lincoln, kcassman1@unl.edu

A. Sheldon Whitney

University of Hawaii

Robert L. Fox

University of Hawaii

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>



Part of the [Plant Sciences Commons](#)

Cassman, Kenneth G.; Whitney, A. Sheldon; and Fox, Robert L., "Symbiotic Nitrogen Fixation by Legumes Requires Extra Phosphorus" (1980). *Agronomy & Horticulture -- Faculty Publications*. 426.
<https://digitalcommons.unl.edu/agronomyfacpub/426>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

ILLUSTRATED CONCEPTS IN TROPICAL AGRICULTURE
A series prepared by the Department of Agronomy and Soil Science
College of Tropical Agriculture and Human Resources
University of Hawaii

SYMBIOTIC NITROGEN FIXATION BY LEGUMES REQUIRES EXTRA PHOSPHORUS



Fig. 1. Soybean supplied with fixed N was scarcely nodulated but developed a vigorous root system.



Fig. 2. Soybean roots that grew in nutrient solution which supplied no N produced abundant nodules and relatively less root mass.

When grain legumes are grown on nitrogen-deficient soils, they may symbiotically fix enough N for good yields, even if no N fertilizer is applied. However, legumes preferentially assimilate N from the soil if this source is available to them. As a result, excess N via the soil may inhibit symbiotic N_2 fixation by legumes.

The physiological and morphological modifications that make symbiotic N_2 fixation by legumes possible include: (1) the invasion of host roots by effective strains of *Rhizobium*, (2) the development of nodules to "house" the *Rhizobium* and, (3) the translocation of mineral nutrients and photosynthate from the host to the nodules to maintain *Rhizobium* activity. Nodules represent an extra sink for plant assimilates, a sink that non-nodulated legumes do not have.

The partitioning of dry matter between roots and nodules is affected by the phosphorus supply. When soybean plants were grown in N-deficient and P-deficient sand culture, nodules comprised 4% of the total plant dry matter and 12% of the root dry weight. When adequate P was supplied in

the nutrient solution, but N was still deficient, nodule dry weight comprised 9% of total plant dry matter and 61% of root dry weight. The P requirement for optimal nodulation and N_2 fixation was greater than the requirement for root or shoot growth. When soybean plants were supplied with combined N in the nutrient solution so that nodule development was poor, root weight and length were greater than in N-fixing plants grown at comparable P levels.

These results suggest that soybean plants that depend upon symbiotic N_2 fixation for their N have a higher requirement for P than their counterparts supplied with fixed N from an external source. A field experiment conducted on Haiku clay soil (Humoxic Tropohumult) showed that N-fixing soybean plants did indeed have a higher external soil P requirement than N-fertilized plants.